Is all salinity the same? I. The effect of ionic compositions on the salinity tolerance of five species of freshwater invertebrates

Zalizniak, Liliana; Kefford, Benjamin; Nugegoda, Dayanthi

https://researchrepository.rmit.edu.au/esploro/outputs/9921859564001341/filesAndLinks?institution=61RMIT_INST&index=null

Zalizniak, L., Kefford, B., & Nugegoda, D. (2006). Is all salinity the same? I. The effect of ionic compositions on the salinity tolerance of five species of freshwater invertebrates. *Marine and Freshwater Research*, 57, 75–82. <https://doi.org/10.1071/mf05103>

Published Version: <https://doi.org/10.1071/mf05103>

Repository homepage: <https://researchrepository.rmit.edu.au>

© CSIRO 2006

Downloaded On 2023/04/26 20:59:45 +1000

**Is all salinity the same? I. The effect of ionic compositions on the salinity
tolerance of five species of freshwater invertebrates.**

Liliana Zalizniak, Ben J. Kefford^A, and Dayanthi Nugegoda

Biotechnology and Environmental Biology, School of Applied Sciences, RMIT

University, PO Box 71, Bundoora 3083, Vic, Australia

^ACorresponding author; email: ben.kefford@rmit.edu.au

Citation:

Zalizniak, L, Kefford, B and Nugegoda, D 2006, 'Is all salinity the same? I. The effect of ionic compositions on the salinity tolerance of five species of freshwater invertebrates', *Marine And Freshwater Research*, vol. 57, pp. 75-82.

Abstract

Salts of marine origin, predominantly consisting of Na^+ and Cl^- ions are dominant in most Australian inland saline waters. The proportions of other ions, Ca^{2+} , Mg^{2+} , SO_4^{2-} , HCO_3^- and CO_3^{2-} , in the water may influence salinity tolerance of freshwater organisms and thus the effect of increasing salinity may vary with difference in ionic proportions. We exposed freshwater invertebrates to different concentrations of four ionic compositions and compared them to the commercial sea salt, Ocean Nature. They were: synthetic Ocean Nature (ONS) and three saline water types (ONS but without [1]: SO_4^{2-} , HCO_3^- and CO_3^{2-} , [2]: Ca^{2+} , HCO_3^- and CO_3^{2-} , [3]: Ca^{2+} , Mg^{2+}) which are considered to be the predominant saline water types in southeastern Australia and the Western Australian wheatbelt. The 96-h LC_{50} values for the five media were determined for six invertebrate species and sub-lethal responses were observed for two species. There were no differences between responses of invertebrates to various ionic compositions in acute toxicity tests. However in prolonged sub-lethal tests animals reacted differently in the various ionic compositions. The greatest effect was observed in water types lacking Ca for which plausible physiological mechanisms exist. Variation in ionic proportions should be taken into account when considering sub-lethal effects of salinity on freshwater invertebrates.

Keywords: salinity, ionic compositions, freshwater invertebrates, toxicity

1 **Introduction**

2

3 The salinization of freshwaters is a major environmental concern in all continents
4 with large arid and semiarid regions, including Australia (Williams 1987). Recently
5 attention has been given to the lethal (Berezina 2003, Kefford *et al.* 2003) and sub-
6 lethal tolerance (Kefford and Nuggeoda 2005a) of freshwater invertebrates to
7 increased salinity, while other studies have experimentally considered effects of
8 salinity on freshwater invertebrate communities (Nielsen *et al.* 2003, Marshall and
9 Bailey 2004). All of these studies used artificial sea salts, the ionic proportion of
10 which approximates seawater, because it is the most common composition of saline
11 water bodies of southeastern Australia (Bayly and Williams 1973), which are sodium
12 chloride (NaCl) dominated. However recently it has been acknowledged that there is
13 some variation in the ionic proportion of NaCl-dominated inland saline waters of
14 southeastern Australia (Radke *et al.* 2002, 2003). The three major saline water types
15 existing in southeastern Australia (Radke *et al.* 2002), and the wheatbelt region of
16 Western Australia (Pinder *et al.* 2005), were proposed by Drever (1982) and occur
17 due to precipitation out of solution of specific minerals during evapoconcentration of
18 saline waters and result in reductions in the relative concentrations of specific ions. If
19 variations in ionic proportions in NaCl-dominated inland saline waters result in
20 differing biological effects, then studies investigating the effects of saline water with a
21 particular ionic proportions (such as seawater) may not accurately describe the effects
22 of changes in salinity with differing ionic proportions. Consequently, we investigated
23 whether these three common ionic proportions and artificial seawater altered lethal
24 and sub-lethal effects of salinity on freshwater invertebrates. For the common ionic
25 proportions we used the most extreme cases where specific ions are eliminated from a

1 saline water source, and therefore refer to the ionic compositions (presence/absence of
2 specific ions), because if the absence of specific ions do not affect salinity tolerance
3 then it is very unlikely that a reduction in the proportions of these ions would affect
4 salinity tolerance.

6 **Materials and methods**

8 ***Test animals***

9 Six species of freshwater invertebrates were used for acute 96-h LC₅₀ toxicity testing
10 (LC₅₀ is the concentration of a toxicant lethal to 50% of a population). The protozoan
11 *Paramecium caudatum* Ehrenberg and hydrozoan *Hydra oligactis* Pallas were
12 purchased from Southern Biological, Nunawading, Victoria, Australia. Other species,
13 collected from central Victoria, in the southern end of the Murray-Darling Basin were:
14 gastropod *Physa acuta* Draparnaud (Campaspe River, at the Kyneton-Heathcote Rd.
15 (37°23'S 144°31'E)), caddis fly *Notalina fulva* Kimmins, water bug *Micronecta*
16 *robusta* Hale and mayfly *Centroptilum* sp. (King Parrot Creek, a tributary of the
17 Goulburn River, at Flowerdale (37°23'S, 145°16'E). These specific species were
18 chosen because they represent a wide range of different taxonomic groups found in
19 freshwaters and were obtainable in sufficient numbers to experimentally expose to
20 varying salinity and ionic composition treatments. Previous experiments where
21 macroinvertebrate species have been collected from different sites or from the same
22 site on different dates have shown no detectable difference in acute lethal salinity
23 tolerance (Kefford *et al.* 2003, 2005, unpublished data). This is despite large
24 differences in the acute lethal salinity tolerance between species. We thus assume that
25 any difference in salinity tolerance or response to the different ionic compositions of

species obtained from different sources represent differences between the species tested rather than differing responses of animals collected from different sites.

Water quality data from collection sites are given in an auxiliary publication, Table 1.

Three species, *P. acuta*, *P. caudatum* and *H. oligactis* were used in chronic toxicity tests. The results for *P. acuta* will be presented elsewhere (Zalizniak *et al.* in prep). For hydra and paramecia the culture growth in different types of treatments was determined as the measure of sub-lethal toxicity and EC₅₀ values calculated (EC₅₀ being the concentration of a toxicant that produced the effect in 50% of population). For *H. oligactis* another sub-lethal end point, tentacle retraction, was used.

Preparation of solutions

Five different solutions were tested. Concentrated stock solution of around 40 mS/cm of Ocean Nature artificial sea salt (ON) (Aquasonic, Wauchope, NSW) was prepared in Milli-Q water and used in preparation of dilutions. Based on both the manufacturers claimed elemental composition and elemental analysis (ICP-MS) of Ocean Nature, 'Ocean Nature Synthesized' (ONS) was prepared from analytical grade reagents. Major ions and trace elements were considered (22 total), and their quantities calculated (see auxiliary publication, Table 2). ON was used as a standard to compare with previous investigations using this salt (Kefford *et al.* 2003, 2004a,b, 2005a,b), and ONS was used as control for possible effects of synthesized various ionic compositions. Based on ONS preparation three different ionic compositions were derived to reproduce the three major saline water types described in Radke *et al.* (2002): S1 had the same ionic composition as ONS except there was no sulphates (SO₄²⁻) and no carbonates (HCO₃⁻ and CO₃²⁻, referred to as alkalinity (Alk)); S2 was

1 without calcium (Ca^{2+}) and Alk, and S3 had Ca^{2+} and magnesium (Mg^{2+}) excluded
2 (Fig. 1; also see auxiliary publication Table 2). Natural S1, S2 and S3 waters have
3 some levels of the elements (see Radke 2002), which we excluded. We excluded them
4 in the stock solutions to represent a worst-case scenario. The control and dilution
5 water had enough of these excluded ions to allow high (>85%) survival. Where
6 possible we tried to use carbon filtered Melbourne tap water (WLW) as our dilution
7 water and control. However, lab cultures required specific media for their
8 maintenance. For paramecia we used Lozina-Lozinsky medium (Lozina-Lozinsky
9 1931), and for hydras – M4 medium (Elendt and Bias 1990). Though M4 medium was
10 designed for daphnids, prolonged culturing of hydra (over several months) using this
11 medium was successful. These media served as culture media, dilution water and
12 control in corresponding experiments. The analysis of major ions for some of these
13 media is presented in auxiliary publication, Table 3.

14 ***Animal cultures***

16 Brown hydra *H. oligactis* was fed daily with brine shrimps or juvenile *Daphnia*
17 *carinata* (whatever available, since previous observations showed that the cultures
18 survive equally well with either) *ad lib*. Medium was replaced three times a week.

19 For paramecia *P. caudatum* culturing technique was per Sazonova *et al.*
20 (1997). Lozina-Lozinsky medium was boiled with 0.4 g L^{-1} of dry yeasts and cooled,
21 and then inoculate of culture was introduced. After two days of acclimatising, animals
22 were used for the experiment. Medium was replaced weekly or as necessary.

23 Animals collected from the field were transported from the site to the
24 laboratory and transferred to the testing solutions as quickly as possible (as per
25 Kefford *et al.* 2003, 2004a, 2005b).

1

2 *Acute toxicity testing experimental protocols*

3 There was no water replacement or feeding.

4

5 *Hydras*

6 The protocol published by Pollino and Holdway (1999) was used and is only briefly
7 described. Non-budding hydras were used. To achieve this hydra were not fed for 1-2
8 days. Five concentrations of each salt type were used: 4, 6, 8, 10 and 12 mS cm⁻¹
9 replicated 4 times in Petri dishes (Ø 54 mm), with 5 animals per replicate and 15 mL
10 of test solution. Observations were made daily for 96 hours; deaths and tentacle
11 retraction of hydras were recorded. For the tentacle retraction only two rankings were
12 used: 'unaffected' being normal, and 'affected', which is any degree of shortening or
13 disintegration (Pollino and Holdway 1999). It is not certain in the tulip stage if
14 animals are truly dead; consequently at the end of experiment tulip stage animals were
15 transferred to control solution for 48 hours. If animals recovered from the tulip stage,
16 they were counted as alive.

17

18 *Paramecia*

19 Five concentrations of each salt type were used: 2, 4, 6, 8, and 10 mS cm⁻¹ each with
20 10 animals per concentration held individually in 2-mL wells. Paramecia were fed
21 with the suspension of yeasts (10 g L⁻¹) in Lozina-Lozinsky medium every second day
22 (0.02 mL per well). Mortality and numbers present was recorded and LC₅₀ was
23 determined after 24, 48 and 96 hours of exposure.

24

25 *Insects*

1 The rapid toxicity testing method was used (Kefford *et al.* 2003, 2005b, in press).
2 There were 10 animals of each species per treatment of 3L of water. Exposure
3 concentrations were: *Centropetillum* sp.: WLW ($EC \approx 0.13 \pm 0.01$ mS cm⁻¹), 5, 10, 15 and
4 20 mS cm⁻¹, and other species: WLW, 10, 15, 20, 25 and 30 mS cm⁻¹. Observations
5 were made daily for 96 hours.

6

7 ***Sub-lethal toxicity testing experimental protocols***

8 *Hydra*

9 Experimental procedure was as per acute test (Pollino and Holdway 1999) and is only
10 briefly described. However, budding hydras were used. To achieve this hydra were
11 fed in excess for 4-6 days. Three concentrations of each salt type were used: 1, 2 and
12 4 mS cm⁻¹. After counting animals and observing tentacle retraction, animals were fed
13 in excess with brine shrimps (0.2 mL per dish). After 1 hour all solutions were
14 changed. All parameters for each day were calculated as the geometrical mean
15 between new and old medium.

16 The mean relative growth rate of hydra for each treatment concentration was
17 calculated as follows (Pollino and Holdway 1999):

18
$$K = (\ln N_t - \ln N_{t-1}) / \Delta t$$

19 Where N_t is the number of animals at time t ,

20 N_{t-1} the number of animals at time of previous observation

21 Δt time between two observations.

22

23 *Paramecia*

24 The experimental protocol is as per acute toxicity testing with paramecia. The culture
25 growth rate (for individual animals) was calculated using a standard formula:

1
$$\mu = \ln N/T$$

2 where N is number of animals in the well at time T

3 and T is time from the start of the experiment (days).

4

5 ***Statistics***

6 For each species and treatment type, Probit regression models (see Agresti 1990) were
7 fitted with the x-variable being EC and the y-variable the response variable (survival,
8 population growth or tentacle contraction). From these regressions LC_{50} and EC_{50}
9 values and their 95% confidence intervals were calculated for each treatment type.
10 Post hoc comparison of EC_{50} values was performed using a paired t-test assuming
11 unequal variances.

12

13 **Results**

14

15 ***Acute tests***

16 For all species examined there were no statistically significant differences in their 96-
17 h LC_{50} values for the different types of treatments (Table 1). The results for
18 *Centroptilum* sp. (Table 1) are, however, somewhat inconclusive. For treatments other
19 than ON over 96 hours of exposure, they had partial but $< 50\%$ mortality at the
20 lowest salinity treatment, 5 mS cm^{-1} , consequently their 96-h LC_{50} value is below 5
21 mS cm^{-1} for all types of treatments except ON. Since concentrations below 5 mS cm^{-1}
22 were not tested in this experiment, the error in LC_{50} calculation is higher than for the
23 other species and thus there is a greater probability of a type 2 error. Across the three
24 species, however, there would appear to be no detectable effect of the different saline
25 water types on acute survival of freshwater invertebrates tested.

Sub-lethal tests

Though there are differences in tentacle retraction of hydra at 24 hours, they were eliminated at 72 hours (Fig. 2). The EC₅₀ for S3 salt type initially increased then later decreased. Thus the hydras appear to adapt to their environment when the initial shock is reduced, and they can return to their 'normal' condition. Interestingly 24- and 48-h EC₅₀ for S1 seemed higher (though not statistically significant) than all the others. It may be that sulphates are more toxic to hydra than chlorides and eliminating them results in a marginally reduced overall toxicity.

Hydra culture growth was partially affected by the variation in ionic compositions (Fig. 3). Ninety six-hour EC₅₀ value for S2 treatment was significantly lower than for the ONS and S3 types of treatments.

The population growth of the paramecia was significantly reduced (Fig. 4), when Ca was eliminated from the media (S2 and S3 types).

Discussion

General observations

There were no significant differences in toxicity between ON, ONS and S1 (no sulphates and alkalinity) treatments in any of the experiments. While it was expected with ON and ONS, it also indicated that removal of SO₄²⁻ and Alk did not change the toxicity of salinity in any detectable way. The proportion of these anions is around 13% of the total anions load in ONS, the rest being mostly Cl⁻. When S1 and S2 treatments were prepared these anions were replaced with Cl⁻, thus increasing its load. Kefford *et al.* (2004a) observed that ON was less toxic to freshwater invertebrates than pure NaCl. The lack of a difference in toxicity between ON, ONS and S1 may

1 indicate that the difference in toxicity between ON and NaCl is not because of Cl⁻
2 toxicity, but rather lack or difficulty in extraction at high salinity of essential and trace
3 elements, such as calcium, potassium, copper, selenium etc. 24- and 48-h EC₅₀ for
4 hydra's tentacle retraction in S1 were slightly higher than in other treatments. We did
5 not specifically test toxicity of Cl⁻ against SO₄²⁻, but other studies with a range of
6 freshwater invertebrate taxa indicate that Na₂SO₄ is more toxic than NaCl (Goetsch
7 and Palmer 1987; Kefford *et al.* 2004a; Palmer *et al.* 2004) and that NaCl is more
8 toxic than ON (Kefford *et al.* 2004a). It would therefore appear that SO₄²⁻ is more
9 toxic than Cl⁻. The replacing of SO₄²⁻ with Cl⁻ could thus have slightly reduced the
10 overall toxicity to hydra.

11 The results regarding treatments with Ca deficiencies are discussed in detail
12 below.

13

14 ***Acute tests***

15 Short-term acute toxicity testing is usually conducted in sub-optimal conditions for
16 animals tested: static water regime and no food supply. Though these tests convey
17 very useful information on the range of tolerance of the animals to a particular
18 toxicant, which can be very useful in modelling and management on a wider scale,
19 they give very little information on the mechanisms of action or the effects of a
20 toxicant to organisms subject to long exposures and low sub-lethal concentrations.
21 These experiments are therefore usually regarded as a starting point for more detailed
22 long-term sub-lethal exposures, from which one can get more definite information on
23 the effects of a particular toxicant. Though both species were clearly affected by the
24 different ionic compositions in our sub-lethal experiments, it was not so in the acute
25 tests (Table 1). In a short-term exposure with lethal concentrations of salinity, the

different ionic compositions had no detectable effect. Osmoregulatory mechanisms may have played a major part in combating the effects of high salinity, rather than fine-tuned biochemical and physiological interactions. Chapman *et al.* (2000) found that there were no differences in the survival or swim-up fry toxicity tests (96-h exposure) of rainbow trout embryos in two saline effluents with different ionic proportions. However they found that chironomid larvae grew differently in the different effluent (10-d exposure). The same results were obtained for sulphates-dominated saline lakes in the USA (Dickerson *et al.* 1996). Though the researchers stated that undiluted lake water was toxic to *Ceriodaphnia dubia* and attributed this to the differences in ionic composition of major ions, when we recalculated LC₅₀ (% of dilution) provided by the authors, the LC₅₀ in terms of electrical conductivities were surprisingly similar and not significantly different for *C. dubia* (except in very saline waters) and fathead minnows. These studies and our results consistently indicate that the short-term lethal toxicity of saline solutions found in nature is not generally affected by different ionic proportion/composition, but longer exposures or sub-lethal effects can reveal the differences. Salinity produced from pure salts (e.g. NaCl, Na₂SO₄) and one to one ratio of pure salts, neither of which occur in nature, however, do have differing toxicity to that of mixtures of salts (Mount *et al.* 1997, Kefford *et al.* 2004a, Palmer *et al.* 2004).

Sub-lethal tests

There could be several explanations regarding the chronic sub-lethal effects of varying ionic compositions:

- (1) Direct effect of deficiency of the essential element Ca.

1 (2) Indirect effect of hardness cations (Ca^{2+} , Mg^{2+}) and carbonates on the
2 biochemistry of the trace-metals.

3

4 ***Direct effects of deficiencies in Ca***

5 *Effects on paramecia*

6 Paramecia have around 5000 cilia. Movement of the cilia is controlled by their
7 membrane potential. Stimulation of cilia (chemically or physically) activates a
8 voltage-sensitive Ca^{2+} current associated with the ciliary membrane (Preston and
9 Hammond 1998). This results in avoidance behaviour, making paramecia swim
10 backward (Preston *et al.* 1992). Nakaoka and Ooi (1985) found that in the presence of
11 ATP as a stimulus in the medium, paramecia swim forward if Ca^{2+} concentration is
12 below 10^{-6} M ($40 \mu\text{g L}^{-1}$) and backward if it is higher than 10^{-6} M. This suggests that,
13 though directional swimming is governed by the *intracellular* Ca^{2+} concentration, a
14 minimum amount of calcium in medium is required to maintain normal responses to
15 stimuli. Slightly proportionally higher concentrations of trace metals in S2 and S3
16 (especially at higher salinities) might have affected animals, but lack of calcium in
17 these media did not allow them to respond adequately. In the case of acute toxicity
18 (Table 1) the differences between various ionic composition types were not evident
19 possibly because short-term effect of higher salinity *per se* was greater than the effect
20 of ionic composition of media, making osmoregulatory mechanism primarily
21 responsible for mortality. At lower salinities in sub-lethal exposures calcium
22 deficiencies might play a greater part in paramecia swimming behaviour, thus making
23 animals in Ca^{2+} -lacking media more prone to abnormal behaviour, and consequently
24 expending more energy. In addition morphogenesis of the complex cell surface
25 during mitosis involves transcellular wave signal, which involves cortical alveoli that

1 act as Ca reservoir in the cell (Laurent and Fleury 1995). Presumably if there were
2 not enough Ca to initiate the signal, mitosis would be abnormal.

3

4 *Effect on hydra*

5 External Ca^{2+} ions play a major role in the nematocyst discharge in hydrozoans
6 (Salleo *et al.* 1994a,b, Yanagita 1973, McKay and Anderson 1988; cited in Kawaii *et*
7 *al.* 1999). Santoro and Salleo (1991) observed that nematocytes do not discharge in
8 Ca^{2+} -free medium, and that La^{3+} , Cd^{2+} and Co^{2+} prevented discharge by blocking the
9 Ca^{2+} channel even when some Ca^{2+} is present. Gitter *et al.* (1994) found that discharge
10 of the stenoteles (a type of nematocyst) in *Hydra vulgaris* is regulated by a
11 mechanism, allowing intake of Ca^{2+} from ambient solution. This may explain why
12 hydras were not affected in the acute toxicity test (involving no feeding and therefore
13 no nematocysts discharge) (Table 1), but were growing slower in sub-lethal test
14 (where nematocysts were discharged to capture their prey) in the S2 treatment
15 compared to ONS control (Fig. 3). As there was some Ca^{2+} present in the M4
16 medium, which was used as control and dilution water for the range of salinities
17 prepared, at higher salinities the effect of blocking Ca^{2+} by increasing concentrations
18 of Co^{2+} and Ni^{2+} (see Auxiliary publication, Table 2) may have begun to play a role.
19 Kawaii *et al.* (1999) reported that Mg^{2+} also had an inhibitory effect on *atrichous*
20 *isorhiza* (a type of nematocyst) discharge, and that the inhibitory effect of Mg^{2+}
21 increased when the external concentration of Ca^{2+} was lowered. This might explain
22 why the S2 type affected sub-lethal salinity tolerance in hydra. S3 type medium,
23 though lacking Ca^{2+} , may not affect hydra as much as an S2 type (Fig. 3) because it
24 also lacked a powerful Ca^{2+} blocker i.e. Mg^{2+} .

Freshwater hydras reproduce by means of forming buds and developing a foot at the base of a bud and then detaching from the parent. A separated bud was counted as a new animal in our experiments. Zeretzke *et al.* (2002) found that in *Hydra vulgaris* (Zurich strain) foot formation was prevented by lowered concentrations of ambient Ca^{2+} , making animals form branches, that persisted on parent's body instead. It would definitely affect the culture growth in our study, as the number of separate individuals has not increased.

Increased toxicity of trace metals

Water quality parameters such as hardness and alkalinity can influence the interactions of ions in the ambient solution. Increases in hardness have shown to result in decreased copper toxicity to fish (Pagenkopf 1983) and cladocerans *Daphnia magna* (Schamphelaere and Janssen 2002) as a result of competition between the hardness metals (Ca, Mg) and trace-metal species for interaction sites. Welsh *et al.* (2000) also showed that acute copper toxicity was lower in waters containing proportionately more Ca. They also indicated that Ca is more important than Mg in modifying the toxicity of Cu in rainbow trout and chinook salmon. The same applies to uptake of zinc by rainbow trout (Alsop and Wood 1999) and *D. magna* (Heijerick *et al.* 2001), cadmium by *D. magna* (Penttinen *et al.* 1998) and the amphipod *Hyalella azteca* (Jackson *et al.* 2000), and manganese by brown trout (Stubblefield *et al.* 1997) in the presence of competing Ca^{2+} ions. All water types used in our study contained essential and trace metals Fe, Mn, Cu, Zn, Mo, Se, Li, Sr, Br, Rb, Co, V, Ni (auxiliary publication, Table 2) that at elevated concentrations can be toxic to aquatic invertebrates. Though the concentration of each trace element was very low, a combined load might be significant in the absence of calcium. Elimination of Ca

1 and/or Mg out of the solution can result, first, in the relative increase of
2 concentrations of trace elements, especially at higher salinities, and second, in
3 increased toxicity of these elements because in the absence of Ca and/or Mg more
4 sites are available for binding at the organism-water interface. The hypothesis of
5 increased trace metals toxicity in Ca^{2+} lacking media remains to be tested.

6 Metal toxicity can also be reduced by complexation with carbonate, thus
7 decreasing the activity of free hydrated metal ions (Barata *et al.* 1998).

9 **Conclusions**

10
11 Variation in ionic compositions common in saline inland waters of southeastern
12 Australia did not affect acute lethal salinity tolerance of any species investigated.
13 However the different ionic compositions affected the three sub-lethal responses of
14 investigated species. The water types lacking Ca had sub-lethally most deleterious
15 effects on the animals. The different responses of invertebrates to various ionic
16 composition types in combination with the sub-lethal range of salinity may be
17 governed by deficiencies of Ca, the chemical interaction of hardness cations,
18 alkalinity and trace metal uptake and toxicity.

19 In assessing the effects of salinity on freshwater invertebrates the ionic
20 proportions should be considered in salinity exposures that are likely to induce sub-
21 lethal effects.

23 **Acknowledgments**

24 We are grateful for funding from Land and Water Australia (LWA) and the Murray
25 Darling Basin Commission, under the National Rivers Contaminants Program (LWA

1 Project no. RMI 12), and the Queensland Department of Natural Resources and
2 Mines. We thank Satish Choy, Brendan Edgar, Richard Marchant, Leon Metzeling,
3 Daryl Nielsen, Carolyn Palmer and Phil Papas for their assistance to the project by
4 being members of a steering committee. We also thank Victor Zalizniak for assistance
5 in calculation of ionic proportions of the media.

References

- Agresti, A. (1990). Categorical data analysis. (John Wiley & Sons, New York, USA)
- Alsop, D.H., and Wood, C.M. (1999). Influence of waterborne cations on zinc uptake and toxicity in rainbow trout, *Oncorhynchus mykiss*. *Canadian Journal of Fisheries and Aquatic Sciences* **56**(11), 2112-2119.
- Barata, C., Baird, D.J., and Markich, S.J. (1998). Influence of genetic and environmental factors on the tolerance of *Daphnia magna* Straus to essential and non-essential metals. *Aquatic Toxicology* **42**, 115-137.
- Bayly, I. A. E., and Williams, W. D. (1973). 'Inland waters and their ecology.' Camberwell, Longman.
- Chapman, P.M., Bailey, H., and Canaria, E. (2000). Toxicity of total dissolved solids associated with two mine effluents to chironomid larvae and early life stages of rainbow trout. *Environmental Toxicology and Chemistry* **19**(1), 210-214.
- Dickerson, K.K., Hubert, W.A., and Bergman, H.L. (1996). Toxicity assessment of water from lakes and wetlands receiving irrigation drain water. *Environmental Toxicology and Chemistry* **15**(7), 1097-1101.
- Drever, J.J. (1982). 'The Geochemistry of Natural Waters.' 2nd Edn. (Prentice-Hall Inc., New Jersey, USA.)
- Elendt, B.-P., and Bias, W.-R. (1990). Trace nutrient deficiency in *Daphnia magna* cultured in standard medium for toxicity testing. Effects of the optimization of culture conditions on life history parameters of *D. magna*. *Water Research* **24**(9), 1157-1167.
- Gitter, A.H., Oliver, D., and Thurm, U. (1994). Calcium- and voltage-dependence of nematocyst discharge in *Hydra vulgaris*. *Journal of Comparative Physiology A* **175**, 115-122.

1 Goetsch, P.A., and Palmer, C.G. (1997). Salinity tolerance of selected
2 macroinvertebrates of the Sabie River, Kruger National Park, South Africa.
3 *Archives of Environmental Contamination and Toxicology* **32**, 32-41.

4 Heijerick, D.G., De Schampelaere, K.A.C., and Janssen, C.R. (2002). Predicting
5 acute Zinc toxicity for *Daphnia magna* as a function of key water chemistry
6 characteristics: development and validation of a biotic ligand model.
7 *Environmental Toxicology and Chemistry* **21**(6), 1309-1315.

8 Jackson, B.P., Lasier, P.J., Miller, W.P., and Winger, P.W. (2000). Effects of calcium,
9 magnesium, and sodium on alleviating cadmium toxicity to *Hyaella azteca*.
10 *Bulletin of Environmental Contamination and Toxicology* **64**, 279-286.

11 Kawai, S., Yamashita, K., Nakai, M., Takahashi, M., and Fusetani, N. (1999).
12 Calcium dependence of settlement and nematocyst discharge in actinulae of the
13 hydroid *Tubularia mesembryanthemum*. *The Biological Bulletin* **196**(1), 45-51.

14 Kefford, B.J., Papas, P.J., and Nuggeoda, D. (2003). Relative salinity tolerance of
15 macroinvertebrates from the Barwon River, Victoria, Australia. *Marine and*
16 *Freshwater Research* **54**, 755-765.

17 Kefford, B.J., Palmer, C.G., Pakhomova, L., and Nuggeoda, D. (2004a). Comparing
18 test systems to measure the salinity tolerance of freshwater invertebrates. *Water*
19 *SA*, **30**(4), 499-506.

20 Kefford, B.J., Papas, P.J., Metzeling, L., and Nuggeoda, D. (2004b). Do laboratory
21 salinity tolerances of freshwater animals correspond with their field salinity?
22 *Environmental Pollution* **129**, 355-362.

23 Kefford, B.J., and Nuggeoda, D. (2005a). No evidence for a critical salinity threshold
24 for growth and reproduction in the freshwater snail *Physa acuta*. *Environmental*
25 *Pollution* **134**, 377-383.

1 Kefford, B.J., Palmer, C.G., and Nuggeoda, D. (2005b). Relative salinity tolerance of
2 freshwater macroinvertebrates from the south-east Eastern Cape, South Africa
3 compared with the Barwon Catchment, Victoria, Australia. *Marine and*
4 *Freshwater Research* **56**, 163-171.

5 Kefford, B.J., Palmer, C.G., Jooste, S., Warne, M. St.J., and Nuggeoda, D. (in press).
6 What is it meant by '95% of species'? An argument for the inclusion of rapid
7 tolerance testing. *Human and Ecological Risk Assessment*.

8 Laurent, M., and Fleury, A. (1995). A model with excitability and relay properties for
9 the generation and the propagation of a Ca^{2+} morphogenetic wave in
10 *Paramecium*. *Journal of Theoretical Biology* **174**(2), 227-236.

11 Lozina-Lozinsky, L.K. (1931). Ernährungsphysiologie der infusorien. *Archiv fur*
12 *Protistenkunde* **74**, 18-120.

13 McKay, M.C., and Anderson, P.A.V. (1988). Preparation and properties of cnidocytes
14 from the sea anemone *Anthopleura elegantissima*. *The Biological Bulletin* **174**,
15 47-53.

16 Marshall, N. A. and P. C. E. Bailey (2004). Impact of secondary salinisation on
17 freshwater ecosystems: effects of contrasting, experimental, short-term releases
18 of saline wastewater on macroinvertebrates in a lowland stream. *Marine and*
19 *Freshwater Research* **55**(5), 509-523.

20 Mount, D.R., Gulley, D.D., Hockett, J.R., Garrison, T.D. and Evans, J.M. (1997).
21 Statistical models to predict the toxicity of major ions to *Ceriodaphnia dubia*,
22 *Daphnia magna* and *Pimephales promelas* (flathead minnows). *Environmental*
23 *Toxicology and Chemistry* **16**, 2009-2019.

- 1 Nakaoka, Y., and Ooi, H. (1985). Regulation of ciliary reversal in triton-extracted
2 Paramecium by calcium and cyclic adenosine monophosphate. *Journal of Cell*
3 *Science* **77**(1), 185-195.
- 4 Nielsen, D. L., M. Brock, Crossle, K., Harris, K., Healey, M., and Jarosinski, I.
5 (2003). The effects of salinity on aquatic plant germination and zooplankton
6 hatching from two wetlands sediments. *Freshwater Biology* **48**, 2214-2223.
- 7 Pagenkopf, G.K. (1983). Gill surface interaction Model for trace-metal toxicity to
8 fishes: Role of complexation, pH, and water hardness. *Environmental Science*
9 *and Technology* **17**, 342-347.
- 10 Palmer, C.G., Muller, W.J., Gordon, A.K., Scherman, P-A, Davies-Coleman, H.,
11 Pakhomova, L. and de Kock, E. (2004). The development of a toxicity database
12 using freshwater macroinvertebrates, and its application to the protection of
13 South African water resources. *South African Journal of Science* **100**, 643-650.
- 14 Pinder, A.M., Halse, S.A., McRae, J.M. and Shiel, R.J. (2005). Occurrence of aquatic
15 invertebrates of the wheatbelt region of Western Australia in relation to salinity.
16 *Hydrobiologia* **543**, 1-24.
- 17 Pollino, C.A., and Holdway, D.A. (1999). Potential of two hydra species as standard
18 toxicity test animals. *Ecotoxicology and Environmental Safety* **43**, 309-316.
- 19 Penttinen, S., Kostamo, A., and Kukkonen, J.V.K. (1998). Combined effects of
20 dissolved organic material and water hardness on toxicity of cadmium to
21 *Daphnia magna*. *Environmental Toxicology and Chemistry* **17**(12), 2498-2503.
- 22 Preston, R.R., Saimi, Y., and Kung, C. (1992). Calcium-dependent inactivation of the
23 calcium current activated upon hyperpolarization of *Paramecium tetraurelia*.
24 *The Journal of General Physiology* **100**, 253-268.

1 Preston, R.R., and Hammond, J.A. (1998). Long-term adaptation of Ca^{2+} -dependent
2 behaviour in *Paramecium tetraurelia*. *Journal of Experimental Biology* **201**,
3 1835-1846.

4 Radke, L.C., Howard, K.W.F., and Gell, P.A. (2002). Chemical diversity in south-
5 eastern Australian saline lakes I: geochemical causes. *Marine and Freshwater*
6 *Research* **53**, 941-959.

7 Radke, L.C., Juggins, S., Halse, S.A., De Deckker, P., and Finston, T. (2003).
8 Chemical diversity in south-eastern Australian saline lakes II: biotic
9 implications. *Marine and Freshwater Research* **54**, 895-912.

10 Salleo, A., La Spada, G., and Barbera, R. (1994a). Gadolinium is a powerful blocker
11 of the activation of nematocytes of *Pelagia noctiluca*. *Journal of Experimental*
12 *Biology* **187**, 201-206.

13 Salleo, A., La Spada, G., Drago, M., and Curcio, G. (1994b). Hyposmotic shock-
14 induced discharge in acontia of *Caliactis parasitica* is blocked by gadolinium.
15 *Experientia* **50**, 148-152.

16 Santoro, G., and Salleo, A. (1991). The discharge of *in situ* nematocysts of the acontia
17 of *Aiptasia mutabilis* is a Ca^{2+} - induced response. *Journal of Experimental*
18 *Biology* **156**, 173-185.

19 Sazonova, V.E., Zaliznyak, L.A., Savel'eva, L.M., Morozova, E.V., and Kostyuk,
20 O.B. (1997). Use of bioassays to develop monitoring of water ecosystem.
21 *Russian Journal of Ecology* **28**(3), 207-212.

22 Schamphelaere, K.A.C. de, and Janssen, C.R. (2002). A biotic ligand model
23 predicting acute copper toxicity for *Daphnia magna*: The effects of calcium,
24 magnesium, Sodium, Potassium, and pH. *Environmental Science and*
25 *Technology* **36**, 48-54.

1 Stubblefield, W.A., Brinkman, S.F., Davies, P.H., Garrison, T.D., Hockett, J.R., and
2 McIntyre, M.W. (1997). Effects of water hardness on the toxicity of manganese
3 to developing brown trout (*Salmo trutta*). *Environmental Toxicology and*
4 *Chemistry* **16**(10), 2082-2089.

5 Williams, W.D. (1987). Salinization of rivers and streams: an important
6 environmental hazard. *AMBIO* **16**(4), 180-185.

7 Welsh, P.G., Lipton, J., Chapman, G.A., and Podrabsky, T.L. (2000). Relative
8 importance of calcium and magnesium in hardness-based modification of copper
9 toxicity. *Environmental Toxicology and Chemistry* **19**(6), 1624-1631.

10 Yanagita, T.M. (1973). The 'cnidoblast' as an excitable system. *Publications of Seto*
11 *Marine Biological Laboratory* **20**, 675-693.

12 Zalizniak, L., Kefford, B.J., and Nugegoda, D. (in prep). Is all salinity the same? II.
13 Effects of different ionic compositions on survival and growth of *Physa acuta*.

14 Zeretzke, S., Perez, F., Velden, K., and Berking, S. (2002). Ca²⁺-ions and pattern
15 control in Hydra. *International Journal of Developmental Biology* **46**(5), 705-71

1 Table 1. The LC₅₀ values for animal species tested in acute 96-h experiments.

Species	Type of treatment	LC ₅₀ values (95% confidence intervals)			
		24-h	48-h	72-h	96-h
<i>P. caudatum</i>	ON	8.70 (7.81-9.67)	8.70 (7.81-9.67)	NM	8.70 (7.81-9.67)
	ONS	8.66 (7.77-9.62)	8.66 (7.77-9.62)	NM	8.66 (7.77-9.62)
	S1	9.10 (8.17-10.17)	8.85 (7.93-9.88)	NM	8.85 (7.93-9.88)
	S2	7.24 (6.24-7.82)	7.24 (6.24-7.82)	NM	7.24 (6.24-7.82)
	S3	7.58 (6.65-8.40)	7.38 (6.47-8.15)	NM	7.38 (6.47-8.15)
<i>H. oligactis</i>	ON	8.95 (8.50-9.48)	8.75 (8.33-9.32)	8.56 (8.15-9.22)	8.37 (7.91-9.21)
	ONS	9.08 (8.60-9.57)	8.79 (8.37-9.30)	8.61 (8.20-9.15)	8.35 (7.90-8.96)
	S1	9.09 (8.63-9.57)	9.09 (8.63-9.57)	8.90 (8.47-9.40)	8.81 (8.39-9.32)
	S2	9.12 (8.66-9.58)	8.86 (8.43-9.32)	8.86 (8.43-9.32)	8.53 (8.07-9.00)
	S3	9.10 (8.63-9.57)	9.10 (8.63-9.57)	8.92 (8.49-9.39)	8.33 (7.76-8.84)
<i>N. fulva</i>	ON	NC	40.55	22.96 (20.32-25.56)	18.46 (16.10-20.94)
	ONS	NC	33.03 (28.91-119.74)	28.17 (24.32-37.90)	19.58 (7.93-23.90)
	S1	NC	32.83	22.55 (19.85-25.55)	18.64 (15.89-21.14)
	S2	60.18	29.51 (26.74-38.27)	24.20	17.97 (14.18-21.54)
	S3	97.46	23.66 (21.28-26.91)	18.27 (15.64-21.04)	15.69 (13.21-17.92)
<i>M. robusta</i>	ON	21.44	19.08	14.51 (-8.27-24.92)	13.44 (-99.73-32.06)
	ONS	25.15 (23.17-27.54)	21.45	18.70	10.51
	S1	29.71 (25.42-45.73)	23.67	17.90	15.78 (2.96-26.14)
	S2	27.61 (25.57-29.67)	23.76	18.02	16.46
	S3	24.15 (15.81-83.88)	19.01	14.11	11.22 (6.65-14.60)
<i>Centropilum</i> sp.	ON	14.94 (12.58-16.97)	9.25	6.60	5.58
	ONS	13.61	6.33 (2.70-9.26)	2.46 (-3.49-5.33)	1.75 (-3.58-4.21)
	S1	14.37 (12.42-16.76)	10.24	6.57	4.63
	S2	11.32 (8.91-13.49)	7.89	5.17	3.79
	S3	10.19 (7.75-12.53)	6.38 (3.71-8.94)	4.11 (0.07-6.89)	3.57 (-0.95-6.36)

2
3 NM – not measured, NC – not calculated (100% survival in all concentrations). For some values CI could not be calculated.

1 a)

2

3 b)

4

5 Fig. 1. Measured ionic proportions of the various saline water types, media (M4) and
6 WLW (see auxiliary publication Table 3 for raw data): a) cations and b) anions as a
7 percentage of the total major cations/anions on a mass to volume basis.

1

2 Fig. 2. Values of EC_{50} (tentacle retraction) for *H. oligactis* in different types of
 3 treatment (error bars indicate 95% CI).

1 Fig. 3. Ninety six-hour EC_{50} values (culture growth) for *H. oligactis* in different types
 2 of treatment (Mean $\pm$ SE, N=4). Different letters represent significantly different
 3 results.

1 Fig.4. Ninety six-hour EC_{50} values (culture growth) for *P. caudatum* in different types
 2 of treatment (Mean $\pm$ SE, N=10). Different letters represent significantly different
 3 results.